



Yilmaz, T. I., Duschl, F., & Di Genova, D. (2016). Feathery and network-like filamentous textures as indicators for the re-crystallization of quartz from a metastable silica precursor at the Rusey Fault Zone, Cornwall, UK. *Solid Earth and Discussions*, 7(6), 1509–1519.
<https://doi.org/10.5194/se-7-1509-2016>

Publisher's PDF, also known as Version of record

License (if available):
CC BY

Link to published version (if available):
[10.5194/se-7-1509-2016](https://doi.org/10.5194/se-7-1509-2016)

[Link to publication record in Explore Bristol Research](#)
PDF-document

This is the final published version of the article (version of record). It first appeared online via EGU at <http://www.solid-earth.net/7/1509/2016/>. Please refer to any applicable terms of use of the publisher.

University of Bristol - Explore Bristol Research

General rights

This document is made available in accordance with publisher policies. Please cite only the published version using the reference above. Full terms of use are available:
<http://www.bristol.ac.uk/red/research-policy/pure/user-guides/ebr-terms/>



Supplement of

Feathery and network-like filamentous textures as indicators for the recrystallization of quartz from a metastable silica precursor at the Rusey Fault Zone, Cornwall, UK

Tim I. Yilmaz et al.

Correspondence to: Tim I. Yilmaz (tim.yilmaz@min.uni-muenchen.de)

The copyright of individual parts of the supplement might differ from the CC-BY 3.0 licence.

Supplementary

5 **Table S1: Compilation of measured concentrations for major and trace elements across the area shown in Fig. 8; C = clear quartz core, FT = feathery texture.**

Texture	FT	C	C	FT
Spot	18	19	20	21
<i>Major elements (wt%)</i>				
SiO ₂	99.53	99.20	99.20	99.63
TiO ₂	0.0013	0.0019	0.0020	0.0012
Al ₂ O ₃	0.473	0.754	0.66	0.358
K ₂ O	0.025	0.003	0.019	0.010
Li ₂ O	0.046	0.136	0.097	0.044
<i>total</i>	<i>100.09</i>	<i>100.09</i>	<i>99.99</i>	<i>100.04</i>
<i>Trace elements (ppm)</i>				
Fe	---	---	---	---
Mn	128	---	95	57
Mg	0.2	---	2.6	0.4
Ca	57	---	---	---
Na	15	2	13	9
P	70	---	61	---
Cu	0.07	0.08	0.06	0.03
Sb	0.5	0.5	0.6	0.29
Pb	---	---	---	---
B	0.1	---	0.8	1.4
Cr	0.11	---	---	---
Ge	3.4	3.4	3.2	1.7
Rb	1.2	0.2	1.0	0.6
U	---	---	---	---




Figure S1: Photomicrograph of subhedral to euhedral quartz crystals from the Rusey fault zone. The two rectangles represent positions of Raman measurements. Further measurements were performed in quartz crystals with different *c*-axis orientations; the arrows indicate three quartz crystals with different *c*-axis orientations; the *c*-axis of the bright crystal is horizontally aligned to the thin section plane; the *c*-axis of the black crystal is vertically aligned to the thin section plane; the *c*-axis of the medium grey quartz crystal has a intermediate alignment.